



Tumor Registry

| Primary Site | Measure Specifications | COC Benchmark Compliance Rate | Methodist Compliance Rate | |
|--------------|---|-------------------------------|---------------------------|------|
| | | | 2013 | 2014 |
| Breast | (NQF #219) Radiation therapy is administered within 1 year (365 days) of diagnosis for women under age 70 receiving breast conserving surgery for breast cancer. (BCSRT) | 90% | 97% | 100% |
| | (NQF #0559) Combination chemotherapy is considered or administered within 4 months (120 days) of diagnosis for women under 70 with AJCC T1cN0M0, or stage II or III hormone receptor negative breast cancer. (MAC) | 90% | 100% | 95% |
| | (NQF #0220) Tamoxifen or third generation aromatase inhibitor is considered or administered within 1 year (365 days) of diagnosis for women with AJCC T1cN0M0, or stage II or III hormone receptor positive breast cancer. (HT) | 90% | 97% | 99% |
| | Radiation therapy is considered or administered following any mastectomy within 1 year (365 days) of diagnosis of breast cancer for women with >4 positive regional lymph nodes. (MASTRT) | 90% | 91% | 100% |
| | (NQF #0221) Image or palpation-guided needle biopsy (core or FNA) is performed to the primary site to establish diagnosis of breast cancer. (nBx) | 80% | 82% | 84% |
| Colon | (NQF #0223) Adjuvant chemotherapy is considered or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC Stage III (lymph node positive) colon cancer. (ACT) | 90% | 100% | 100% |
| | (NQF #0225) At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer. (12RLN) | 85% | 95% | 85% |
| Gastric | At least 15 regional lymph nodes are removed and pathologically examined for resected gastric cancer. (G15RLN) | NA | 57% | 0% |
| Lung | Surgery is not the first course of treatment for cN2, M0 lung cases. (LNoSurg) | NA | 100% | 100% |
| | Systemic chemotherapy is administered within 4 months to the day preoperatively or day of surgery to 6 months postoperatively or it is considered for surgically resected cases with pathologic lymph node - positive (pN1 and pN2) NSCLC. (LCT) | NA | 100% | 100% |
| Rectum | Preoperative chemo and radiation are administered for clinical AJCC T3N0, T4N0, or Stage III; or Postoperative chemo and radiation are administered within 180 days of diagnosis for clinical AJCC T1-2N0 with pathologic AJCC T3N0, T4N0, or Stage III; or treatment is considered; for patients under the age of 80 receiving resection for rectal cancer. (RECRCT) | NA | 100% | 100% |